



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : C07D 307/18, 405/10, 407/10	A1	(11) International Publication Number: WO 89/ 02892 (43) International Publication Date: 6 April 1989 (06.04.89)
---	-----------	---

(21) International Application Number: PCT/US88/03347
(22) International Filing Date: 28 September 1988 (28.09.88)
(31) Priority Application Number: 103,484
(32) Priority Date: 30 September 1987 (30.09.87)
(33) Priority Country: US

(71) Applicant: THE UNIVERSITY OF TENNESSEE RE-
SEARCH CORPORATION [US/US]; Suite 415,
Communications Building, Knoxville, TN 37996
(US).

(72) Inventors: KENNEDY, Thomas, P. ; 1624 Clanlo
Drive, Memphis, TN 38104 (US). KABALKA,
George, W. ; 7124 Cresthill Drive, Knoxville, TN
37919 (US).

(74) Agents: HODGES, Paul, E. et al.; Luedeka, Hodges &
Neely, 1030 First American Center, Knoxville, TN
37902 (US).

(81) Designated States: AT (European patent), AU, BE (Eu-
ropean patent), CH (European patent), DE (Euro-
pean patent), FR (European patent), GB (European
patent), IT (European patent), JP, LU (European pa-
tent), NL (European patent), SE (European patent).


Published

With international search report
With amended claims.

Date of publication of the amended claims:

20 April 1989 (20.04.89)

(54) Title: CERTAIN 3-SUBSTITUTED 2-ALKYL BENZOFURAN DERIVATIVES

**(57) Abstract**

The disclosure relates to compounds of formula (I) and pharmaceutically acceptable addition salts thereof wherein X represents a single, direct bond or a substituted or unsubstituted alkylene chain containing 1 to 4 carbon atoms, wherein R₅ is a lower alkyl group, wherein R₆ is either hydrogen or methyl, wherein Am is selected from the class consisting of amino, lower mono and dialkylamino, piperidino, piperazino, N-lower alkyl piperazino, pyrrolidino, and morpholino groups wherein Y₁ and Y₂ are identical and are hydrogen, a halogen, methyl or ethyl and n is an integer in the range of 1-5. They are useful in treating arrhythmic conditions and conditions for which treatment with a vasodilator is indicated.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

AMENDED CLAIMS

[received by the International Bureau on 30 March 1989 (30.03.89);
original claim 1 amended; other claims unchanged (1 page)]

Claim 1. A compound of the formula:



10

and pharmaceutically acceptable addition salts thereof where-
in X represents a single, direct bond or a substituted or
unsubstituted alkylene chain containing 1 to 4 carbon atoms
where such substituents are one or more members selected from
15 the group consisting of branched or straight-chain alkyl,

cycloalkyl, aryl, alkoxy and the formula -O-C(=O)-R_4 , with R_4
being hydrogen or lower alkyl, wherein R_5 is a lower alkyl
group, wherein R_6 is either hydrogen or methyl, wherein Am is
20 a group selected from the class consisting of amino, lower
mono and dialkylamino, piperidino, piperazino, N-lower alkyl
piperazino, pyrrolidino, and morpholino groups, wherein Y_1
and Y_2 are identical and are selected from the class consist-
ing of hydrogen, halogen, methyl and ethyl and n is an inte-
25 ger in the range of 1-5.

2. A compound as set forth in Claim 1

wherein X represents an alkylene chain having the formula -C(=O)-
wherein R_1 and R_2 are each selected from the class consisting
5 of hydrogen, a group having the formula -OR_3 with R_3 being a

lower alkyl group, and a group having the formula -O-C(=O)-R_4
with R_4 being hydrogen or a lower alkyl group, R_4 is butyl,
10 R_5 is hydrogen, Am is selected from the class consisting of
amino and lower mono and dialkylamino and Y_1 and Y_2 are
identical and are selected from the class consisting of



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : C07D 307/18, 405/10, 407/10	A1	(11) International Publication Number: WO 89/ 02892 (43) International Publication Date: 6 April 1989 (06.04.89)
(21) International Application Number: PCT/US88/03347 (22) International Filing Date: 28 September 1988 (28.09.88) (31) Priority Application Number: 103,484 (32) Priority Date: 30 September 1987 (30.09.87) (33) Priority Country: US (71) Applicant: THE UNIVERSITY OF TENNESSEE RESEARCH CORPORATION [US/US]; Suite 415, Communications Building, Knoxville, TN 37996 (US). (72) Inventors: KENNEDY, Thomas, P. ; 1624 Clanlo Drive, Memphis, TN 38104 (US). KABALKA, George, W. ; 7124 Cresthill Drive, Knoxville, TN 37919 (US).		(74) Agents: HODGES, Paul, E. et al.; Luedeka, Hodges & Neely, 1030 First American Center, Knoxville, TN 37902 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: CERTAIN 3-SUBSTITUTED 2-ALKYL BENZOFURAN DERIVATIVES <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div> (57) Abstract <p>The disclosure relates to compounds of formula (I) and pharmaceutically acceptable addition salts thereof wherein X represents a single, direct bond or a substituted or unsubstituted alkylene chain containing 1 to 4 carbon atoms, wherein R₅ is a lower alkyl group, wherein R₆ is either hydrogen or methyl, wherein Am is selected from the class consisting of amino, lower mono and dialkylamino, piperidino, piperazino, N-lower alkyl piperazino, pyrrolidino, and morpholino groups wherein Y₁ and Y₂ are identical and are hydrogen, a halogen, methyl or ethyl and n is an integer in the range of 1-5. They are useful in treating arrhythmic conditions and conditions for which treatment with a vasodilator is indicated.</p>		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

CERTAIN 3-SUBSTITUTED 2-ALKYL BENZOFURAN DERIVATIVES

The invention relates to compounds having pharmacological activity and more particularly relates to novel pharmacologically active 3-substituted 2-alkyl benzofuran derivatives, and methods for their preparation.

5 Compounds in accordance with the invention are represented by the general formula:



15 and pharmaceutically acceptable addition salts thereof wherein X represents a single, direct bond or a substituted or unsubstituted alkylene chain containing 1 to 4 carbon atoms, wherein R₅ is a lower alkyl group, wherein R₆ is either hydrogen or methyl, wherein Am is a group selected
20 from the class consisting of amino, lower mono and dialkylamino, piperidino, piperizino, N-lower alkyl piperizino, pyrrolidino, and morpholino groups, wherein Y₁ and Y₂ are identical and are hydrogen, halogen, methyl or ethyl, and n is an integer in the range of 1-5.

25 The term "unsubstituted or substituted alkylene chain containing 1 to 4 carbon atoms" is intended, unless further defined, to designate a saturated aliphatic hydrocarbon chain of between 1 and 4 carbon atoms with or without one or more substituents. Substituents are limited
30 to those which do not diminish the pharmacological activity of the compounds below a useful level and include branched or straight-chain alkyl or cycloalkyl groups, aryl groups,

alkoxy groups, and ester substituents. "Lower alkyl" is intended to designate straight-chain, branched, or cyclic saturated aliphatic hydrocarbon groups containing 1-6 carbon atoms. "Lower mono and dialkylamino" refers to amino groups with one or two straight-chain, branched or cyclic saturated aliphatic hydrocarbon groups containing 1-6 carbon atoms. When two groups are present, they may be the same or different. Examples are methylamino, dimethylamino, ethylamino, diethylamino, n-propylamino, isopropylamino, and the like. Halogen, unless further defined, is intended to refer to fluorine, chlorine, bromine, and iodine.

Compounds in accordance with the invention are useful as vasodilators and as antiarrhythmic agents. Preferred for this purpose are compounds of Formula I above

wherein X represents the Formula $\begin{array}{c} R_1 \\ | \\ -C- \\ | \\ R_2 \end{array}$ above wherein R_1 and/or

R_2 are hydrogen, lower alkyl groups, groups with the Formula $-OR_3$ with R_3 being a lower alkyl group, or groups with the

Formula $\begin{array}{c} O \\ || \\ -O-C-R_4 \end{array}$ with R_4 being hydrogen or a lower alkyl group, R_5 is a lower alkyl group containing 1-4 carbon atoms, R_6 is hydrogen, Am is as defined above for Formula I, Y_1 and Y_2 are identical and are hydrogen, bromine, iodine, or methyl, and n is in the range of 1-3. Particularly preferred

are compounds wherein X is $\begin{array}{c} R_1 \\ | \\ -C- \\ | \\ R_2 \end{array}$ wherein R_2 is hydrogen and R_1

is hydrogen, or $-OR_3$ with R_3 being a lower alkyl group, or R_1


is $\begin{array}{c} O \\ || \\ -O-C-R_4 \end{array}$ with R_4 being hydrogen or a lower alkyl group, R_5 is butyl, R_6 is hydrogen, Am is amino or lower mono and dialkyl amino, Y_1 and Y_2 are identical and are hydrogen, bromine, iodine, or methyl and n is an integer in the range

of 1-3. Most preferably, X is $\begin{array}{c} R_1 \\ | \\ -C- \\ | \\ R_2 \end{array}$ wherein R_2 is hydrogen and

R_1 is hydrogen or $-OR_3$ with R_3 being a lower alkyl group containing between 1 and 4 carbon atoms, or R_1 is $-O-\overset{\overset{O}{\parallel}}{C}-R_4$ with R_4 being hydrogen or a lower alkyl group containing 1 to 4 carbon atoms, R_5 is n-butyl, R_6 is hydrogen, Am is amino, ethylamino or diethylamino, Y_1 and Y_2 are either both hydrogen, both iodine, or both methyl, and n is 1. Of the most preferred compounds, compounds where R_1 and R_2 are both hydrogen are particularly desirable.

10 Compounds of Formula I in which R_6 is hydrogen are prepared by first condensing an alkali metal salt of a compound represented by Formula II below in which X, R_5 , Y_1 and Y_2 have the same meanings as in Formula I with a
15 dibromoalkane represented by Formula III in which R_6 is hydrogen and n is 1-5 in an inert organic medium such as dimethyl formamide.


20



II


25

30



III

The resulting bromoalkoxy-substituted compounds of Formula IV are condensed with an amine of the Formula V in which Am has
35 the same meaning as in Formula I in an inert solvent such as benzene to produce the Formula I compounds.




H-Am

V

10 Alternately, when Am does not represent a secondary amine and R_6 is either hydrogen or methyl, an alkali metal salt of a compound of Formula II can be condensed with an amine represented by Formula VI in which Z is a halogen atom to produce of Formula I compounds.


15



20


The compounds represented by Formula II can be synthesized by a number of reaction routes. As will become more apparent hereinafter, many of such compounds can be prepared by reduction of or reduction and subsequent reaction of a ketone intermediate represented by Formula VII wherein A is a single direct bond or a substituted or unsubstituted alkalene chain containing 1-3 carbon atoms in the chain and R_5 , Y_1 , and Y_2 are as defined in Formula I.

30




When A is a single, direct bond, Formula VII ketones are known intermediates and are disclosed in U.S. Patent Nos. 3,248,401 and 3,920,707, which are incorporated herein by reference.

5 When A represents a substituted or unsubstituted alkalene chain containing 0-3 carbon atoms, the ketone intermediates represented by Formula VII can be prepared by Friedel-Crafts acylation of a 2, 6-substituted anisole of
10 Formula IX with an acid chloride of Formula VIII wherein m represents an integer of 0-3 and R_7 and R_8 represent the same entities as R_1 and R_2 or precursors thereof followed by demethylation of the anisole with pyridine hydrochloride.






VIII





IX

The acid chlorides of Formula VIII can be prepared from 3-carboxy-2-alkyl benzofurans of the Formula X by reaction with
35 an alkene Grignard reagent of Formula XI wherein o is 0-2 and R_7 and R_8 are defined as in Formula VIII in the presence of


CdCl_2 to result in the formation of the secondary alcohols of Formula XII.



Formula XII alcohols can be dehydrated to the corresponding alkenes of Formula XIII below by reaction with sulfonyl chloride in pyridine followed by reaction with lithium triethyl borohydride. Formula XIII alkene substituted benzofuran compounds are converted to acid chlorides of Formula VIII by ozonation in the presence of zinc and oxidation of the resulting aldehyde of Formula XIV to the carboxylic acid employing potassium permanganate (cold) followed by reaction with sulfonyl chloride.




15 When X is a single direct bond, compounds of
 Formula I are prepared by first reacting a compound of
 Formula XV with Y_1 and Y_2 as defined in Formula I with ethyl
 bromoacetate in acetone in the presence of K_2CO_3 to form a
 compound of Formula XVI. Formula XVI compounds are converted
 20 to the compounds of Formula XVII by Perkin condensation in
 the presence of acetic anhydride and sodium acetate followed
 by the conversion of the ester group to R_5 groups.
 Demethylation of the substituted anisole moiety yields
 compounds of Formula II wherein X is a single, direct bond
 25 which can be employed as previously discussed to prepare
 Formula I compounds.




-8-

5



XVI

10




XVII

15 Compounds of Formula XV can be prepared by Friedel-Crafts acylation of the substituted anisoles of Formula IX with salicyloyl chloride.

The particularly preferred compounds of Formula I
 20 described above wherein X is $\begin{array}{c} R_1 \\ | \\ -C- \\ | \\ R_2 \end{array}$ and R₂ is hydrogen and R₂ is

hydrogen or -OR₃ with R₃ being a lower alkyl group or $-O-\overset{\overset{O}{\parallel}}{C}-R_4$
 25 with R₄ being hydrogen or a lower alkyl group are advantageously prepared by way of an alcohol intermediate which is produced by reducing a ketone of the formula:


30



XVIII

35

with R_4 , R_5 , R_6 , Y_1 and Y_2 , and n as defined for Formula I. Formula XVII ketones are known and procedures for their synthesis are described in U.S. Patent Nos. 3,248,401 and 3,920,707, the disclosures of which are incorporated herein by reference. To produce compounds according to Formula I wherein Y_1 and Y_2 are identical halogens, reduction of the compounds of Formula XVIII with Y_1 and Y_2 being halogens is performed under conditions which reduce the ketone group to the alcohol without otherwise affecting the molecule. A reducing system employing sodium borohydride in a tetrahydrofuran-methanol mixture (10:1 v/v) at approximately 0°C produces high yields of the alcohol represented by Formula XIX:




To prepare compounds of the invention wherein Y_1 and Y_2 are both hydrogen, both methyl, or both ethyl, the ketones of the Formula XIV wherein Y_1 and Y_2 are both hydrogen, both methyl or both ethyl are similarly reduced to produce the alcohol intermediate shown in Formula XX. Alternately, to produce the compounds where Y_1 and Y_2 are both hydrogen, reduction of Formula XVIII compounds wherein Y_1 and Y_2 are both halogens can be performed employing a reduction system which reduces the ketone group to the alcohol while also dehalogenating the benzene ring to produce Formula XX alcohols. Sodium borohydride in methanol in the presence of a $PdCl_2$ catalyst at 20°C is a preferred reduction

-10-

system to achieve both reduction and dehalogenation.

5



XX


10

15

Compounds of Formula I wherein X is $\begin{matrix} R_1 \\ | \\ -C- \\ | \\ R_2 \end{matrix}$ and R_1 (and


R_2) is hydrogen are produced from the intermediates of Formulas XIX and XX by further reduction at the alcohol group. Compounds of Formula XXI (Y_1 and Y_2 are both halogens, methyl or ethyl) or XXII (Y_1 and Y_2 are both hydrogen), when reacted in a suitable solvent at 0°C with sodium borohydride in trifluoroacetic acid produce compounds of Formulas XXI and XXII, respectively.

25



XXI

30



XXII


35

-11-


The alcohols of Formulas XIX and XX are also employed as intermediates to produce compounds wherein X is

$\begin{array}{c} R_1 \\ | \\ C \\ | \\ R_2 \end{array}$ and R_2 is hydrogen and R_1 is $-OR_3$ with R_3 being a lower alkyl group. A Williamson synthesis whereby the alcohols or Formula XIX or XVII are converted to the corresponding alkoxide and reacted with an alkyl halide of the Formula R_3X is used to produce the ethers represented by Formulas XXIII

10 (Y_1 and Y_2 are both halogens, methyl or ethyl) and XXIV (Y_1 and Y_2 are both hydrogen).



XXIII




XXIV

To produce the compounds of Formula I wherein X is

$\begin{array}{c} R_1 \\ | \\ -C- \\ | \\ R_2 \end{array}$ and R_1 is $-O-\overset{\overset{O}{\parallel}}{C}-R_4$ (R_2 is hydrogen), the alcohols of

5 Formulas XIX and XX are esterified. Acyl halides of the
 formula $R_4-\overset{\overset{O}{\parallel}}{C}-X$ can be reacted with the alcohols of Formulas
 XIX or XX, respectively, preferably in the presence of a
 solvent capable of acting as an acid scavenger, e.g.,
 10 pyridine, to produce compounds of Formulas XXV (halogenated)
 (Y_1 and Y_2 are both halogen, methyl or ethyl) or XXVI (Y_1 and
 Y_2 are both hydrogen), respectively:


15



XXV

20

25



XXVI

30

The compounds of Formula I react to form acid
 addition salts with pharmaceutically acceptable acids, for
 35 example, with inorganic acids, such as hydrochloric acid,
 hydrobromic acid, sulphuric acid and phosphoric acid and with

organic acids such as acetic acid, tartaric acid, maleic acid, citric acid and toluenesulfonic acid.

The compounds of the Formula I above and the salts thereof are useful in treating arrhythmic conditions and conditions for which treatment with a vasodilator is indicated. The novel pharmaceutically active agents provided by the present invention can be administered in pharmaceutical dosage forms, internally, for example, parenterally or enterally with dosage adjusted to fit the exigencies of the therapeutic situation. The pharmaceutical dosage forms are prepared by incorporating the active ingredient in conventional liquid or solid vehicles to thereby provide emulsions, suspensions, tablets, capsules, powders and the like according to acceptable pharmaceutical practices. A wide variety of carriers or diluents as well as emulsifying agents, dispersing agents and other pharmaceutically acceptable adjuvants can be incorporated in the pharmaceutical dosage forms.

The following examples are offered to illustrate the invention and are not intended to be limiting.

EXAMPLE I

Preparation of (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl methanol.

One 1 mmole (645 mg) of the ketone (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl - methanone is dissolved in 30 ml of THF:MeOH (10:1 v/v). Sodium borohydride (1.2 mmole, 45.42 mg) is added to the solution and the mixture is stirred and maintained at a temperature of 0°C until the starting material is consumed (15 minutes). Excess borohydride is destroyed by the dropwise addition of water (0.5 ml). Volatile components are removed under reduced pressure (roto-evaporator). Water is added to the residue (10 ml) followed by the addition of methylene chloride (10 ml). The methylene chloride layer is separated from the aqueous phase and is dried over anhydrous sodium sulfate. The methylene chloride solvent is removed

under reduced pressure and the product is purified by column chromatography (silica gel support using methylene chloride) and is recovered by reduced pressure evaporation of the methylene chloride. The yield of the product, m.p. 106-5 107°C, is >50% of theoretical. (The m.p. of the hydrochloride salt is 143-145°C.)

EXAMPLE II

10 Preparation of (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-phenyl methanol.

One mmole (645 mg) of the ketone, (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl methanone is dissolved in 10 ml of methanol. Palladium
15 dichloride (2 mmole, 354 mg) is added and the mixture is agitated to suspend the palladium dichloride. The temperature of the mixture is adjusted to 20°C. Sodium borohydride (10 mmole, 379 mg) is added and stirring is continued until reaction is complete (1 hour). The
20 palladium dichloride is removed by filtration and water is added to the filtrate. An ether extraction is performed and the product is removed from the ether phase by evaporation under reduced pressure. The produce is purified by chromatography (silica gel using methylene chloride) and
25 results in >50% yield of the product, m.p. 203°C (decomposes).

EXAMPLE III

Preparation of (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl methane.

30 One mmole (647 mg) of the alcohol as prepared in EXAMPLE I is dissolved in methylene chloride (5 ml). Sodium borohydride (38 mg, 10 mmole) added to 10 ml of trifluoroacetic acid and the mixture is cooled to 0°C. The methylene chloride solution is added slowly to the
35 trifluoroacetic acid solution and the mixture stirred for 30 minutes at 0°C. Excess borohydride is destroyed by the

dropwise addition of water (0.5 ml). Volatile components are removed under reduced pressure (roto-evaporator). Water is added to the residue (25 ml) followed by the addition of methylene chloride (25 ml). The methylene chloride layer is separated, washed twice with 25 ml of 5% aqueous sodium hydroxide and 25 ml of water. The methylene chloride solution is dried over sodium sulphate and then passed through a short (5 cm) basic alumina column. Evaporation of the solvent yields the product, m.p. 80-81°C, in >70% yield.
10 (The m.p. of the hydrochloride salt is 119-121°C.)

EXAMPLE IV

Preparation of methoxy (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl methane.

One mmole (647 mg) of the alcohol as prepared in
15 EXAMPLE I is dissolved in 10 ml of THF. The solution is cooled to -78°C and lithium diisopropylamide in cyclohexane (1.1 mmole, 0.73 ml of a 1.5 M solution) is slowly added. Methyl iodide (1.2 mmole, 0.17 g) is added and the mixture permitted to warm to room temperature (30 minutes). The
20 volatile components are removed under reduced pressure (roto-evaporator) and the residue is dissolved in methylene chloride. The methylene chloride solution is dried over anhydrous sodium sulfate and is purified by passing the solution through silica gel column as in EXAMPLE I. The
21 product, m.p. 96-98°C, is obtained upon evaporation of the solvent in a theoretical yield of >90%.

EXAMPLE V


Preparation of (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl methyl pivalate.
25

One mmole (647 mg) of the alcohol as prepared in
EXAMPLE I is dissolved in pyridine (4 ml). Excess pivaloyl chloride (5 mmole, 605 mg) is added to the pyridine solution and the mixture heated to 65°C until the starting alcohol is
30 completely consumed (approximately 12 hours). Volatile materials are removed under reduced pressure (roto-evaporator). The residue is dissolved in methylene chloride

and the methylene chloride solution washed twice with 25 ml of 5% aqueous sodium hydroxide and once with 25 ml of water. The methylene chloride solution is dried over sodium sulfate and then passed through a short (5 cm) basic alumina column. 5 Evaporation of the solvent yields the product in >90% yield. (The m.p. of the hydrochloride salt is 108-110°C.)

THE CLAIMS:

1. A compound of the formula:



and pharmaceutically acceptable addition salts thereof
 15 wherein X represents a single, direct bond or a substituted
 or unsubstituted alkylene chain containing 1 to 4 carbon
 atoms wherein R_5 is a lower alkyl group, wherein R_6 is either
 hydrogen or methyl, wherein Am is a group selected from the
 class consisting of amino, lower mono and dialkylamino,
 20 piperidino, piperazino, N-lower alkyl piperazino,
 pyrrolidino, and morpholino groups, wherein Y_1 and Y_2 are
 identical and are selected from the class consisting of
 hydrogen, halogen, methyl and ethyl and n is an integer in
 the range of 1-5.

2. A compound as set forth in Claim 1

wherein X represents an alkylene chain having the formula $\begin{matrix} R_1 \\ | \\ -C- \\ | \\ R_2 \end{matrix}$

5 wherein R_1 and R_2 are each selected from the class consisting
 of hydrogen, a group having the formula $-OR_3$ with R_3 being a
 lower alkyl group, and a group having the formula $-O-\overset{\overset{O}{||}}{C}-R_4$
 with R_4 being hydrogen or a lower alkyl group, R_4 is butyl,
 10 R_5 is hydrogen, Am is selected from the class consisting of
 amino and lower mono and dialkylamino and Y_1 and Y_2 are
 identical and are selected from the class consisting of

hydrogen, bromine, iodine, and methyl and n is an integer in the range of 1-3.

3. A compound as set forth in Claim 2 wherein R₂ is hydrogen and R₁ is selected from the class consisting of hydrogen, a group having the formula -OR₃ with R₃ being a

5 lower alkyl group, and a group having the formula $\text{-O}-\overset{\text{O}}{\parallel}\text{C}-\text{R}_4$ with R₄ being hydrogen or a lower alkyl group, R₅ is butyl, R₆ is hydrogen, Am is selected from the class consisting of amino, ethylamino, and dialkylamino, Y₁ and Y₂ are identical and are selected from the class consisting of hydrogen,
10 iodine, and methyl and n is an integer in the range of 1-3.

4. A compound as set forth in Claim 2 wherein R₂ is hydrogen and R₁ is selected from the class consisting of hydrogen, -OR₃ with R₃ being a lower alkyl group containing

5 between 1 and 4 carbon atoms, $\text{-O}-\overset{\text{O}}{\parallel}\text{C}-\text{R}_4$ with R₄ being hydrogen or a lower alkyl containing 1-4 carbon atoms, R₅ is n-butyl, R₆ is hydrogen, Am is amino, ethylamino or diethylamino, Y₁ and Y₂ are identical and are selected from the class consisting of hydrogen, iodine, and methyl, and n
10 is 1.

5. A compound as set forth in Claim 4 wherein both R₁ and R₂ are hydrogen.

6. A compound according to Claim 1 wherein said compound is 2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5 diiodophenyl] methane.

7. A compound according to Claim 1 wherein said compound is (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-phenyl methane.

8. A compound according to Claim 1 wherein said compound is methoxy (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl methane.

9. A compound according to Claim 1 wherein said compound is methoxy (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-phenyl methane.

10. A compound according to Claim 1 wherein said compound is (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-3,5-diiodophenyl methyl pivalate.

11. A compound according to Claim 1 wherein said compound is (2-n-butyl-3-benzofuranyl) 4-[2-(diethylamino)ethoxyl]-phenyl methyl pivalate.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US88/03347

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶ According to International Patent Classification (IPC) or to both National Classification and IPC IPC(4): C07D 307/18; 405/10; 407/10 U.S.C1.: 544/153,376; 546/196; 548/525; 549/469,471											
II. FIELDS SEARCHED <div style="text-align: center; margin-top: 5px;">Minimum Documentation Searched ⁷</div> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%;">Classification System</th> <th style="width: 80%;">Classification Symbols</th> </tr> <tr> <td style="padding: 5px;">U.S.</td> <td style="padding: 5px;">544/153,376; 546/196; 548/525; 549/469,471</td> </tr> </table> <div style="text-align: center; margin-top: 5px;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸</div>			Classification System	Classification Symbols	U.S.	544/153,376; 546/196; 548/525; 549/469,471					
Classification System	Classification Symbols										
U.S.	544/153,376; 546/196; 548/525; 549/469,471										
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹ <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%;">Category [*]</th> <th style="width: 60%;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="width: 30%;">Relevant to Claim No. ¹³</th> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">U.S. , A, 3,248,401 (TONDEUR ET AL.) 26 April 1966. See the entire document.</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1 & 2</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">Y</td> <td style="padding: 5px;">L. FIESER and M. FIESER "Advanced Organic Chemistry" published 1961, by Reinhold (New York), see pages 441-443 especially page 443.</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1 & 2</td> </tr> </table>			Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	X	U.S. , A, 3,248,401 (TONDEUR ET AL.) 26 April 1966. See the entire document.	1 & 2	Y	L. FIESER and M. FIESER "Advanced Organic Chemistry" published 1961, by Reinhold (New York), see pages 441-443 especially page 443.	1 & 2
Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³									
X	U.S. , A, 3,248,401 (TONDEUR ET AL.) 26 April 1966. See the entire document.	1 & 2									
Y	L. FIESER and M. FIESER "Advanced Organic Chemistry" published 1961, by Reinhold (New York), see pages 441-443 especially page 443.	1 & 2									
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>[*] Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>											
IV. CERTIFICATION <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 5px;"> Date of the Actual Completion of the International Search 23 DECEMBER 1988 International Searching Authority ISA/US </td> <td style="width: 50%; padding: 5px;"> Date of Mailing of this International Search Report <div style="font-size: 1.5em; font-weight: bold;">10 FEB 1989</div> Signature of Authorized Officer BERNARD DENTZ </td> </tr> </table>			Date of the Actual Completion of the International Search 23 DECEMBER 1988 International Searching Authority ISA/US	Date of Mailing of this International Search Report <div style="font-size: 1.5em; font-weight: bold;">10 FEB 1989</div> Signature of Authorized Officer BERNARD DENTZ							
Date of the Actual Completion of the International Search 23 DECEMBER 1988 International Searching Authority ISA/US	Date of Mailing of this International Search Report <div style="font-size: 1.5em; font-weight: bold;">10 FEB 1989</div> Signature of Authorized Officer BERNARD DENTZ										

